

Nervous System

White Shaker Disease Syndrome

Lindsey Buracker, DVM and John Robertson, VMD, PhD

Introduction and Overview

White Shaker Disease Syndrome is a neurologic disease seen primarily in dogs with white coats, particularly in West Highland White Terriers, Maltese Terriers (Bagley et al., 1993), and Samoyeds (Cummings et al., 1986). Affected dogs have a very unique generalized tremor (unintentional, rhythmic muscle movements) and typically are 5 months to 3 years old when the disease is first recognized (Yamaya et al., 2004). There is no gender predilection for this disease. Because this disease has been identified in breeds of dogs lacking a white coat (Dachshund, among others; Yamaya et al., 2004), and in the adult years of their life, the terms “shaker dog syndrome”, “white shaker disease” and “little white shaker dog” commonly are used (Yamaya et al., 2004). The cause of the disease is not known and there is little research being done on this condition in any breed.

Because there are several neurologic and neuromuscular diseases that can produce similar clinical signs, it is essential that Westie owners immediately get a thorough evaluation of their dog and that an accurate diagnosis is made.

Symptoms and Diagnosis

Affected dogs have a history of a relatively sudden onset of constant tremors over the entire body, including the head and eyes. These tremors occur when opposing muscle groups alternately contract and relax in a repetitive manner (Smith and Thacker, 2004). Uncontrolled eye movements, referred to as opsoclonus, consist of rapid, involuntary, multidirectional (horizontal and vertical) movements of the eyes.

The tremors are exaggerated by excitement, handling, forced locomotion, and high levels of stress (Summers et al., 1995). Although some affected dogs may have constant tremors, they remain alert and responsive to their owners and environment. These dogs generally retain normal sensory and muscle functions, which are controlled by the cranial nerves. Consequently, they are able to sense when their faces are touched, and their pupils dilate and constrict appropriately in response to changes in light. In some instances, tremors may be severe enough to cause a wobbly, uncoordinated gait, or overreaching with the legs when walking forward (Smith and Thacker, 2004). This latter change in gait is called hypermetria.

Because other diseases can manifest as tremors, they must be ruled out before an appropriate treatment can be administered (Smith and Thacker, 2004). Examples of these diseases include various inflammatory or infectious diseases of the nervous system, epilepsy, and exposure to toxic substances such as moldy food, lead or organophosphates. These possibilities are ruled out by performing an electrophysiological evaluation of nerve function and microscopic examination of a portion of a nerve obtained by biopsy. After a variety of diagnostic tests are performed to eliminate the aforementioned possibilities, dogs with typical clinical signs that lack evidence of other diseases are diagnosed with White Shaker Disease Syndrome.

Occasionally, affected dogs may have a head tilt. Because head tilts also can occur with other central nervous system disease, peripheral neurologic disorders and even with ear problems, these other causes must first be ruled out before a diagnosis of White Shaker Disease Syndrome is made (Smith and Thacker, 2004).

The diagnosis of White Shaker Disease Syndrome is generally made based on the dog’s history, age at onset, and symptoms. Blood cytology, chemistry and x-rays, as well as physical

Common Clinical Findings

Terriers at Increased Risk

5 Months - 3 Years Old

Tremors

examination findings, are usually normal and have not proven valuable to aid in a diagnosis. A sample of cerebrospinal fluid may be collected for analysis, as an increase in the number of lymphocytes has been noted in some cases (Smith and Thacker, 2004). Fortunately, White Shaker Disease Syndrome is rarely a fatal disease.

Krabbe Disease (Globoid Cell Leucodystrophy)

Young Westies also can develop tremors as a result of another neurologic disease that does not appear to be related to White Shaker Disease Syndrome. This disease, which is known as Krabbe disease or globoid cell leucodystrophy (GCL), is a neurologic disease in which a substance that is toxic to the myelin-forming cells in the nervous system accumulates inside nerve cells. Myelin, which makes up most of the white matter in the central nervous system and is present in the peripheral nervous system, is essential for normal nerve function. The accumulation of this substance results in the breakdown of myelin, which leads to severe neurological symptoms such as progressive blindness, seizures, and eventually death.

This disease, which occurs as the result of a deficiency in a specific enzyme called galactocerebrosidase (GALC) that is critical for cellular metabolism, is called a “storage disease.” Deficiency of GALC has been demonstrated not only in the brain but also the liver and kidneys of affected dogs (Fletcher et al., 1972; Yunis et al., 1976, Wenger et al., 1999).

West Highland White and Cairn terriers are the two breeds most affected by GCL, which is inherited as an autosomal

“ The tremors are exaggerated by excitement, handling, forced locomotion, and high levels of stress. ”

recessive trait (Parker et al., 1995; Wenger et al., 1999). Clinical signs typically become evident beginning around 3 months of age, and include ataxia of the hindlimbs, muscle wasting, head and body tremors, and even blindness. In many dogs, there are substantial degenerative changes in the peripheral and autonomic nervous systems (Summers et al., 1995), and affected dogs may die from the disease in less than a year or may be euthanized due to poor quality of life (Parker et al., 1995). Pathologists have identified gray discolored areas of the brain in affected dogs, firmness of the cerebral cortex, and dilation of the ventricles, reflecting tissue loss (Summers et al., 1995). The brains of affected dogs appear smaller than normal, with a notable decrease in the amount of white matter (Summers et al., 1995).

To determine if a dog is affected with GCL, a blood sample is analyzed by polymerase chain reaction (PCR) to identify the enzyme deficiency underlying the condition (Cifti et al., 2000). Although not done commonly, examination of a sample of a nerve obtained by biopsy using electron microscopy may aid in reaching a diagnosis and MRI can also be useful (Cozzi et al., 1998; Wenger et al., 1999).




Figure 1 - One of the most commonly affected areas of the brain in dogs with White Shaker Disease Syndrome is the cerebellum, which is responsible for 'fine control' of movements.

Krabbe disease has been reported in humans, dogs, mice, monkeys, and sheep (Fletcher et al., 1972). Although death usually occurs before affected infants reach 2 years of age, the disease also has been identified in older people. Canine GCL most closely resembles the late-onset, ultimately fatal form of Krabbe disease occurring in human patients.

Causes

Though the cause remains unknown, White Shaker Disease Syndrome is most often associated with mild inflammation of the central nervous system (non-suppurative encephalomyelitis) (Smith and Thacker, 2004). The cerebellum, which is the part of the central nervous system responsible for 'fine control' of movements, is commonly affected. As a result, dysfunction of this part of the brain could be one of the initiators for the tremor. It is not known if the inflammation is the true underlying cause or if there is an associated neurotransmitter abnormality in affected dogs. Further research should be done to rule out that possibility as well as any virus that might serve as the cause of the disease. There has also been some speculation that White Shaker Disease Syndrome can be congenital in some breeds (West Highland White Terriers, Maltese Terriers, and Samoyeds).

Treatment and Prevention

Early diagnosis of the disease is beneficial in treating affected dogs, as many will respond in a few days to immunosuppressive levels of corticosteroids that have anti-inflammatory effects (Yamaya et al., 2004). The tremors can be reduced with

diazepam (Valium), which is used to diminish anxiety or modify behavior, as a muscle relaxant, or an anticonvulsant (Smith and Thacker, 2004). In some cases, dogs will have to remain on a low dose of corticosteroids for the duration of their life in order to remain free of signs of the disorder.

There are a few adverse affects that can occur from taking high doses of corticosteroids. Some of these include vomiting, gastrointestinal bleeding, ulcers, and diarrhea (Smith and Thacker, 2004). Even though these complications can be serious, most can be managed with appropriate care. Unfortunately, there is no known way to prevent the disease.

Some dogs experiencing tremors may have convulsions, and may refuse to eat or seem to be disconnected from their environment just before onset of the seizures. Affected dogs may need to be encouraged to eat and drink. Some owners have noted that hand feeding and raising food and water bowls off the floor is helpful (Swingle C, 2008), and that symptoms can lessen or resolve when the dog is relaxed or sleeping (Summers et al., 1995). Some dogs respond well to being crated in a minimally dark room that is quiet during times of high stress.

In summary, White Shaker Disease Syndrome is a disease that affects primarily white-coated dog breeds, including Westies. Clinical signs, including involuntary tremor, are seen in young dogs. An accurate diagnosis is essential in order to appropriately manage affected dogs, which can be sustained with treatment. Further research is needed to determine the cause(s) of the disease.

Current Research About White Shaker Disease Syndrome

As a result of its low prevalence, there has not been a lot of research on White Shaker Disease Syndrome osteopathy in dogs in recent years. However, there have been a few recent reports regarding the disease and Krabbe disease in people that deserve specific mention.

Cantuti-Castelvetri L, Maravilla E, Marshall M, Tamayo T, D'auria L, Monge J, Jeffries J, Sural-Fehr T, Lopez-Rosas A, Li G, Garcia K, van Breemen R, Vite C, Garcia J, Bongarzone ER. Mechanism of neuromuscular dysfunction in Krabbe disease. *J Neurosci.* 2015 Jan 28;35(4):1606-16.

Krabbe disease in people results in the loss of skeletal muscle mass and function, the cause of which remains unknown. This study was performed to study skeletal muscle cells and the junctions between nerves and muscles (i.e., neuromuscular junctions) from mice and dogs with GLD. The investigators determined that muscular dysfunction in Krabbe disease is compounded by dysfunction of the neuromuscular junctions, and low levels of a specific cellular pathway that is involved in growth of muscle fibers under normal conditions. The investigators propose that in the future it may be possible to either activate this pathway or replace missing proteins with gene therapy to improve the lives of people with this disease. As it is likely that studies such as this will initially be performed in affected dogs, findings from those studies may be important in veterinary practice as well.

Fletcher JL, Williamson P, Horan D, Taylor RM. Clinical signs and neuropathologic abnormalities in working Australian Kelpies with globoid cell leukodystrophy (Krabbe disease). *J Am Vet Med Assoc.* 2010 Sep 15;237(6):682-8.

In 2010, Fletcher and colleagues studied globoid cell leukodystrophy in a group of Australian Kelpies that were deficient in the enzyme that underlies the condition. Their primary goal in this study was to determine whether or not the degree of demyelination and inflammation in the affected dogs' central nervous system correlated with the clinical signs exhibited by the dogs (i.e., tremors and abnormal gait). They reported that the age of onset and rapid progression of the disease, and the clinical signs were corresponded with those seen in West Highland White and Cairn Terriers, and that the clinical signs reflected abnormalities of the cerebellum. Because Australian Kelpies originated from Collie breeds, the investigators concluded that the disease in these dogs may not be due to the same mutations that cause the disease in West Highland White Terriers, Cairn Terriers or Irish Setters.



Acknowledgements

Mr. Matthew Crotts, a medical illustrator in Educational Resources in the College of Veterinary Medicine at the University of Georgia, created the illustration used in this chapter.

Relevant References

- Bagley R, Kornegay J, Wheeler S, Plumer S, Cauzinille L. Generalized tremors in Maltese: clinical findings in seven cases. *J Amer Anim Hosp Assoc* 29: 141-145, 1993
- Cantuti-Castelvetri L, Maravilla E, Marshall M, Tamayo T, D'auria L, Monge J, Jeffries J, Sural-Fehr T, Lopez-Rosas A, Li G, Garcia K, van Breemen R, Vite C, Garcia J, Bongarzone ER. Mechanism of neuromuscular dysfunction in Krabbe disease. *J Neurosci*. 2015 Jan 28;35(4):1606-16.
- Capucchio MT, Prunotto M, Lotti D, Valazza A, Galloni M, Dore B, Pregel P, Amedeo S, Catalano D, Cornaglia E, Schiffer D. Krabbe's disease in two West Highland White terriers. *Clin Neuropathol*. 2008 Sep-Oct;27(5):295-301.
- Ciftci K, Trovitch P. *Applications of genetic engineering in veterinary medicine*, Elsevier Science, Philadelphia, 2000
- Cozzi F, Vite C, Wenger D, Victoria T, Haskins M. MRI and electrophysiological abnormalities in a case of canine globoid cell leucodystrophy, *Journ Small Anim Pract* 39: 401-405, 1998.
- Cummings J, Summers B, DelaHunta A, Lawson C. Tremors in Samoyed pups with oligodendrocyte deficiencies and hypomyelination. *Acta Neuropath* 71: 267-277, 1986
- Fletcher JL, Williamson P, Horan D, Taylor RM. Clinical signs and neuropathologic abnormalities in working Australian Kelpies with globoid cell leukodystrophy (Krabbe disease). *J Am Vet Med Assoc*. 2010 Sep 15;237(6):682-8.
- Fletcher T, Kurtz H. Animal model for human disease: Globoid cell leukodystrophy. *Am J Path* 66: 375-378, 1972
- Graziano AC, Cardile V. History, genetic, and recent advances on Krabbe disease. *Gene*. 2015 Jan 15;555(1):2-13.
- McGraw RA, Carmichael KP. Molecular basis of globoid cell leukodystrophy in Irish setters. *Vet J*. 2006 Mar;171(2):370-2.
- National Institute of Neurological Disorders and Stroke "NINDS Tremor Information Page/Tremor Fact Sheet"
<http://www.ninds.nih.gov/disorders/tremor/tremor.htm>
- Parker A. Little white shakers syndrome: generalized, sporadic, acquired, idiopathic tremors in adult dogs. In J.D. Bonaguara and R.W. Kirk (eds) In J.D. Bonaguara and R.W. Kirk (eds) *Kirk's Current Veterinary Therapy XII Small Animal Practice*. pp. 1126-1127. W.B. Saunders Co., Toronto. 1995
- Smith K, Thacker L. Generalized temors: Identifying a White Shaker dog. <http://www.addl.purdue.edu/newsletters/2004/spring/tremors.htm>
- Summers B, Cummings J, de Lahunta A. Degenerative diseases of the central nervous system. in *Veterinary Neuropathology*, Mosby-Year book, St. Louis, MO, 1995
- Swingle C. "White Shakers Syndrome", Westie Club of America, <http://www.westieclubamerica.com/health/whiteshaker.html>.
- Wenger D, Victoria T, Rafi M, Luzi P, Vanier M, Vite C, Patterson D, Haskins M. Globoid cell leukodystrophy in Cairn and West Highland White Terriers. *J heredit* 90: 138-142, 1999
- Yamaya Y, Iwakami E, Goto M, Koie, H, Watari, T, Tanaka, S, Takeuchi, A, Tokuriki, M. A case of Shaker Dog Disease in a Miniature Dachshund. *J Vet Med Sci* 66: 1159-1160, 2004
- Yunis E, Lee R. The morphologic similarities of human and canine leukodystrophy. *Am J Path* 85: 99-114, 1976